

12-1-2021

The role of chemokines in vertebrate physiology and disease: a concise review

Hebatallah Mahgoub

Department of Pathology, Faculty of Veterinary medicine, Mansoura University,
hebamahgoub@mans.edu.eg

Follow this and additional works at: <https://mvmj.researchcommons.org/home>

How to Cite This Article

Mahgoub, Hebatallah (2021) "The role of chemokines in vertebrate physiology and disease: a concise review," *Mansoura Veterinary Medical Journal*: Vol. 22: Iss. 4, Article 3.

DOI: <https://doi.org/10.21608/mvmj.2021.46848.1014>

This Review Article is brought to you for free and open access by Mansoura Veterinary Medical Journal. It has been accepted for inclusion in Mansoura Veterinary Medical Journal by an authorized editor of Mansoura Veterinary Medical Journal.

The role of chemokines in vertebrate physiology and disease: a concise review



Hebatallah A. Mahgoub

Department of Pathology, Faculty of Veterinary medicine, Mansoura University, Mansoura, 35516, Egypt.

ARTICLE HISTORY

Received: November 3, 2020

Revised: February 15, 2021

Accepted: February 16, 2021

Corresponding author: Hebatallah A.

MahgoubTel: +20112316644

Email: heba.a.mahgoub@gmail.com

ABSTRACT

Chemokines are classified as proteins with chemoattractant activities that have multiple crucial roles in health and disease, where they participate in the processes of development, angiogenesis, hematopoiesis and tumor metastasis. Multiple cells are involved in the production of chemokines. However, the main cells involved in such role are blood monocytes, macrophages and polymorphonuclear leukocytes, where they exert their role in the inflammatory reaction following antigen recognition by tissue phagocytes. Due to their important role in the establishment of successful immune response, several microbes are incriminated in the production of proteins that mimic chemokines. In addition, their receptors could be used by microbes as a portal of entry to host cells, e.g., human immunodeficiency virus. Chemokines showed significant involvement in the pathogenesis of multiple diseases, e.g., thyroid autoimmune diseases, Behçet's disease and atherosclerosis. Presented is a concise minireview on some of the documented roles of chemokines in several physiological and pathological conditions.

Keywords: Chemokines; chemokines receptors; Behçet's disease; thyroid autoimmune disease; cancer

1. Structure and classification

Chemokines can be defined as chemoattractant proteins, secreted by cells to act on other cells, either by attracting them to a specific site or changing their biological behavior [1, 2]. According to their amino acid structure, they are subdivided into either CXCL (16 members), CCL (28 members), XCL (2 members) or CX3CL (one member) chemokines [2]. They serve important roles in multiple physiological and pathological disciplines, development, angiogenesis, hematopoiesis and tumor metastasis [3–10].

2. Chemokine-secreting and chemokine-responsive cells

The main cells involved in chemokine release are blood monocytes, macrophages and polymorphonuclear leukocytes (PMN) with destined activity on PMN (CXCL1, 2, 3, 4, 5, 6, 7, 8, 12 and 15, CCL1, 2, 3, 4, 5, 7, 8, 13, 15, 24 and 24 and CX3CL1), fibroblasts (CXCL1, 2, 3, 4 and 7 and CCL26), endothelial cells (CXCL4, 5, 6, 7, 8, 8, 10, 11, 15 and 16 and CCL22), T cells (CXCL8, 9, 10, 11, 12, 13 and 14, CCL1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 27 and 28, XCL1 and 2 and CX3CL1), monocytes (CXCL12 and 14, CCL1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 13, 14 and 15 and CX3CL1), macrophages (CXCL12, CCL3, 4, 5, 25 and CX3CL1), natural killer T cells (CXCL16), dendritic cells (CXCL12 and 13, CCL2, 3, 4, 5, 7, 8, 13, 15, 16, 17, 18, 19, 20, 21, 22, 25, XCL1 and 2 and CX3CL1), natural killer cells (CXCL10, 11 and 12, CCL2, 3, 4, 5, 6, 7, 8, 16, 21 and 22, XCL1 and 2 and CX3CL1) and B cells (CXCL10, 11, 12, 13 and 14 and CCL4, 6, 12, 18, 19, 20, 21 and 27) [2, 11]. The main regulatory pathway of chemokine release is the inflammatory cascade [2, 12]. Their function in angiogenesis, the initial primordial

germ cell migration in the early embryo, trigeminal neuron migration and the attraction of hematopoietic stem cells to their bone marrow niche are exerted through CXCL12 [10]. They also serve in lymphoid tissue development [8]. Among the cells associated with chemokine function, platelets express both chemokines, CXCL4, β -thromboglobulins, CCL5 and CCL17, and chemokine receptors, CCR (CCR1, 2, 3, 4, 5, 6, 7, 8, 9, and 10) and CXCR (CXCR1, 2, 3, 4, and 5), on their outer cell membrane [13, 14].

3. Chemokines role in B cell responses

CCL28, a β -chemokine, is expressed by mucosal epithelial cells, showing high expression in the salivary gland, small intestine and colon, is specialized in the recruitment of IgA+ plasma cells to mucosal sites [15–19]. CCL28 expression and the associated B cell recruitment was also recorded in the mammary gland [16].

4. Chemokines function in wound healing

Wound healing regularly occurs at the end of the inflammatory process, following the eradication of the causative agent of cellular injury and/or noxious stimuli [20, 21]. In the preliminary stages of the inflammatory process, chemokines play a crucial role in directing leukocytes to the inflammatory site [1, 22], and hence guaranteeing the establishment of an inflammatory process. Chemokines are also essential for the closure of wounds [23], where CX3CR1, a receptor for CX3CL1 (also known as fractalkine), disruption caused a reduction of alpha-smooth muscle actin (a marker for myofibroblasts) and collagen deposition in skin wounds and reduced neovascularization. Similar effects on defective reepithelization were recorded in models of deficient CXCR2 [24]. In addition, CXCL chemokine interleukin (IL)-8 stimulates

keratinocyte migration and proliferation in cutaneous injury [22, 25–28].

5. Chemokines and cancer

CXCL1 expression was detected in gastric, colonic, colorectal, and urinary bladder cancer patients, and it was associated with the expression of vascular endothelial growth factor (VEGF) and phospho-signal transducer and activator of transcription 3 (p-STAT3) in gastric cancers, stimulators for angiogenesis [29–34]. Its expression was an indication of an advanced cancer grade and it increased xenograft tumor growth with its microvasculature. An *in vitro* application of CXCL1 in gastric cancer cell line increased cellular migration and increased the VEGF signaling (which was suppressed by CXCR2 blockage) and p-STAT3 expression. In a different work by Xu et al. [35], gastric cancer cells stimulated CXCL1 release from lymphatic endothelial cells in a co-culture system, which in turn stimulated lymphatic endothelial cell migration and tube formation. In addition, CXCL1/MGSA/GRO-1 (melanoma growth stimulatory activity/growth regulated oncogene) is involved in the mutagenesis or transformation of melanocytes and ovarian cells [36, 37]. CXCL1 was also expressed by ovarian cancer cell lines and increased their invasiveness [38]. Pancreatic ductal adenocarcinoma cells showed high expression of CXCL1 which had a role in tumor progression [39].

Murine CXCL2, expressed by oral squamous cell carcinoma, was incriminated in osteoclast activation and the resulting bone destruction in cases of oral cancers [40]. In addition, CXCL1 and CXCL2 produced from bone marrow adipocytes were incriminated in the osteolysis associated with prostate cancers [41]. Inhibition of CXCL1 and CXCL2 reduced the metastasis of breast and prostate cancers [42–44]. Mammary adenocarcinoma induced the expression of CCL2, CCL5, and CXCL2 chemokines and CCR1, CCR2, CCR3 and CXCR2 chemokine receptors in splenic murine T lymphocytes [45]. Murine CXCL2 was found to be involved in the translocation of calreticulin (surface marker on cancer cells to be identified and uptaken by phagocytes/antigen-presenting cells) to the outer leaflet of the plasma membrane of cancer cells [46–48].

6. Viral mimicry of chemokines, their receptors, or their binding proteins

Besides mimicry of cytokines, some viruses encode proteins that mimic chemokines (e.g. human immunodeficiency virus, respiratory syncytial virus, members of betaherpesvirus and gammaherpesvirus genera, and molluscum contagiosum virus), chemokine receptors (e.g. members of betaherpesvirus and gammaherpesvirus genera and members of poxviridae), or chemokine-binding proteins/chemokine scavengers (e.g. members of alphaherpesvirus, betaherpesvirus, and gammaherpesvirus genera and members of poxviridae) aiming towards modulating the immune response against the invading viral pathogen [49–62].

7. Chemokine receptors acting as a portal of entry for human immunodeficiency virus (HIV) infection

CD4 is considered the primary receptor for HIV, where gp 120 and gp 41 (envelope proteins) mediate virus binding to the target cell membrane followed by consequent interaction with chemokine receptors CCR3 and 5 CCR2B or CXCR4 leading to merging of the host cell membrane and HIV membrane [63–72]. Other chemokines, CCR2, 3, 8 and 9, STRL33, Gpr15, Gpr1, APJ, ChemR23 and CX3CR1, are also used by HIV for host cell entry during *in vitro* infection [64, 73–76]. In addition, highly exposed persistently seronegative individuals showed upregulated expression of chemokines, RANTES (Regulated upon activation, normal T Cell expressed and presumably secreted), MIP-1 α (macrophage inflammatory protein) and MIP-1 β when exposed to *gag* peptide from HIV [77].

8. Chemokine signaling in skin diseases

Being one of the most common skin diseases, psoriasis involves chronic dermatitis evidenced by leukocytic infiltration (T cells, neutrophils and macrophages) of the dermis and the subcutaneous tissue [78]. The most common chemokines incriminated in the recorded leukocytic recruitment are CXCL8 (IL-8) and 10, CX3CL1 (fractalkine), CCL2/MCP-1 (monocyte chemoattractant protein-1), 5 (RANTES), 20 (MIP-3 α), 26 (eotaxin2) and 27 [79, 80].

9. Chemokines and arterial disease

9.1. Hypertension

Being a serious risk factor for cardiovascular disease, a chemokine-mediated immune response is incriminated in the established disease pathogenesis, including MCP-1/CCL2, IP-10/CXCL10 (interferon (IFN)- γ inducible protein), IL-8/CXCL8, RANTES/CCL5, fractalkine (CX3CL1) and their receptors CCR2, CCR5, CXCR1, CXCR2, CXCR3 and CX3CR1, acting mainly through the activation of macrophages and monocytes migration to the vascular wall and causing endothelial cell dysfunction and vascular smooth muscle cells proliferation [81–83].

9.2. Atherosclerosis

Being one of the established complications of arterial hypertension, chemokines also play important role in the pathogenesis of atherosclerosis through the induced IL-8, fractalkine, MCP-1 which cause vascular inflammation as indicated by the observed vascular leukocyte infiltration and endothelial cell dysfunction [81]. Plasmacytoid dendritic cells seem to play important role in the established pathogenesis through the production of IFN- α cytokine and chemokines (CXCL1, CXCL10) [84].

10. Thyroid autoimmune diseases

Chemokine, particularly CXCL9, 10, and 11, and chemokine receptors, CXCR3, mediate T cell responses that play important role in the pathogenesis of thyroid autoimmune

disorders, e.g. Graves' disease [85–89]. Other chemokines, e.g. CCL2 and CCL5 showed variable levels of expression [90].

11. Alzheimer's disease

Chemokines play an eminent role in the physiology of the nervous system [91]. Upon antigenic stimulation, neuroglial cells, endothelial cells, and neurons of the central nervous system, as well as Schwann cells of the peripheral nervous system can release chemokines. In Alzheimer's disease, a neurodegenerative disease caused by the extracellular deposition of amyloid protein, intracellular neurofibrillary tangles, and the lack of neurons and synapses, inflammatory chemokines may play a vital role in its occurrence and development [20, 21, 91]. They mostly induce immune cell migration to the inflammatory site and regulate the migration of neuroglial cells, neurons and neural progenitors to sites of nervous tissue inflammation (neuroinflammation) [91]. MCP-1/CCL2 and its receptor CCR2 (in the serum, the cerebrospinal fluid (CSF) and the brain tissue) are considered as markers for Alzheimer's disease progression and their levels were positively correlated with a rapid loss of cognition and developing dementia. IL-8 also showed a positive correlation with Alzheimer's disease progression in the serum, and the CSF [92]. IP-10/CXCL10 showed an increase in the CSF of Alzheimer's disease patients [93, 94]. In contrast, RANTES/CCL5 showed downregulation in the serum and upregulation in the brain tissue of Alzheimer's disease patients [95, 96]. On the other hand, CXCL12/SDF-1 (stromal cell-derived factor 1) was reduced in the serum and CSF of reported cases of Alzheimer's disease [97].

12. Brain injury

Chemokines are known to be expressed in the vasculature of the blood-brain barrier, e.g. CXCL12, CCL19, CCL20, CCL21 and CCL27, through which endothelial cells regulate leukocytes entry into the central nervous system (CNS), hence called homeostatic chemokines [98, 99]. They also sustain signals for migration, proliferation, or differentiation to neurons and glia cells [100]. In addition, CX3CL1 and CXCL14 are expressed by the CNS and it appears that they are mainly expressed by the neurons [99, 101–103]. CXCL12 appears to be also involved in the process of neurogenesis and neuronal survival, a function also shared by CX3CL1 chemokine [99, 100, 104, 105].

13. Hepatic disease

Upon liver injury or establishment of hepatic disease, activated Kupffer cells secrete immunoregulatory protein including CXCL1, 2, and 8 (IL-8) as chemoattractants for neutrophils to exert their role in the eradication of the injurious agent that usually culminates in the organ healing upon successful eradication [2, 20, 21, 106–109]. Chemokines, CXCL9, CXCL10 and CXCL11, also play fundamental roles in the recruitment of lymphocytes to sites of chronic hepatic inflammation [107, 109, 110].

14. Pulmonary disease

Chemokines, CXCL1, 2, 5 and 8 were revealed to participate in the pathogenesis of acute lung injury through urgent leukocyte recruitment to the inflammatory site [111]. On the other hand, CCL2, CXCL1, 2, 5, 7, 8, 9, 10, 11 and 16, with their respective receptors, seem to play important role in the pathogenesis of the chronic obstructive pulmonary disease, not only causing recruitment to innate immune cells, neutrophils and monocytes but also enhancing the migration of T cells and B cells to target areas of pulmonary inflammation [112–114]. In addition, it was revealed that the activation of chemokine receptors, CXCR3 on T helper 1 cells and CCR4 and 8 and CXCR4 on T helper 2 cells, through their respective chemokine ligands is responsible for the establishment of idiopathic pulmonary fibrosis [115]. Other chemokines, Eotaxin, Eotaxin-2, RANTES, MCP-3, MCP-1, MIP-1 α and CCR3 have roles in allergic airway hyper-reactivity [116].

15. Diabetes

Being one of the leading reasons for chronic renal failure, diabetes culminates in a set of renal degenerative changes that include glomerular hypertrophy, thickening of basement membranes of renal tubules and glomeruli through the accumulation of extracellular matrix leading to renal fibrosis [117–119]. One of the genetic response towards diabetes-associated hyperglycemia is the enhanced transcription of CCL2 chemokines and other modulators of inflammation [120]. In addition, it was revealed that MCP-1 plays a crucial role in the diabetes-associated nephropathy through its known function of macrophage activation [121].

16. Host-microbe homeostasis

One of the recorded models of dysbiosis or disruption in the homeostasis between host and microbes is periodontitis, where regulatory T cells are involved, through the chemoattractant CCL22, in orchestrating the resulting immune response following antigen recognition to exert a balance between immune response and tolerance, controlling inflammation and helping in the establishment of repair mechanism [122–124]. Such regulatory function of regulatory T cells prevents the resulting tissue pathology if their function is otherwise prohibited.

17. Behçet's disease

Behçet's disease, a chronic recurrent systemic inflammatory disorder, is associated with multiple organ pathologies, oral and genital ulcerations, skin lesions, and uveitis [20, 21]. Multiple etiologies were implicated in the disease pathogenesis, microbial antigens, environmental causes, endothelial cell dysfunction, genetic susceptibility, and immunological aberrations [125]. T helper 1 cells play a crucial role in the pathogenesis of Behçet's disease [126, 127]. Also, the expression of T helper 1-related chemokine receptors seems to be crucial in the established pathogenesis

CCR5 and CXCR3 [128–130], where CXCR3 is considered a marker for IFN- γ -producing T cell population.

Conclusion

This review elucidated the roles of chemokines as chemoattractant proteins in various physiologic and pathologic states. They have protective roles in the blood-brain barrier and share in the process of neurogenesis and neuronal survival in CNS. On the other hand, chemokines served in the pathogenesis of the chronic obstructive pulmonary disease, Behçet's disease, thyroid autoimmune disease and chronic hepatic disease through recruitment of T cells. They also shared other cell receptors as viral receptors for HIV.

REFERENCES

- Murphy K, Travers P, Walport M, Janeway C (2012) Janeway's immunobiology, 8th ed. Garland Science; London: Taylor & Francis [distributor], London
- Murphy K, Weaver C (2016) Janeway's Immunobiology, 9th Edition. Garland Science. <https://doi.org/10.1201/9781315533247>
- Koch AE, Polverini PJ, Kunkel SL, Harlow LA, DiPietro LA, Elner VM, Elner SG, Strieter RM (1992) Interleukin-8 as a macrophage-derived mediator of angiogenesis. *Science* 258:1798-1801 . <https://doi.org/10.1126/science.1281554>
- Nagasawa T, Hirota S, Tachibana K, Takakura N, Nishikawa S, Kitamura Y, Yoshida N, Kikutani H, Kishimoto T (1996) Defects of B-cell lymphopoiesis and bone-marrow myelopoiesis in mice lacking the CXCR chemokine receptor. *Nature* 382:635-638 . <https://doi.org/10.1038/382635a0>
- Zou YR, Kottmann AH, Kuroda M, Taniuchi I, Littman DR (1998) Function of the chemokine receptor CXCR4 in hematopoiesis and in cerebellar development. *Nature* 393:595-599 . <https://doi.org/10.1038/31269>
- Martin TA (2001) Interleukin-8 and Angiogenesis. In: Snyder CR, Jiang WG, Matsumoto K, Nakamura T (eds) Growth Factors and their Receptors in Cancer Metastasis. Springer Netherlands, Dordrecht, pp 51-65. https://doi.org/10.1007/0-306-48399-8_3
- Müller A, Homey B, Soto H, Ge N, Catron D, Buchanan ME, McClanahan T, Murphy E, Yuan W, Wagner SN, Barrera JL, Mohar A, Verástegui E, Zlotnik A (2001) Involvement of chemokine receptors in breast cancer metastasis. *Nature* 410:50-56 . <https://doi.org/10.1038/35065016>
- Chensue SW (2001) Molecular Machinations: Chemokine Signals in Host-Pathogen Interactions. *Clin Microbiol Rev* 14:821-835 . <https://doi.org/10.1128/CMR.14.4.821-835.2001>
- Doitsidou M, Reichman-Fried M, Stebler J, Köprunner M, Dörries J, Meyer D, Eguerra CV, Leung T, Raz E (2002) Guidance of Primordial Germ Cell Migration by the Chemokine SDF-1. *Cell* 111:647-659 . [https://doi.org/10.1016/S0092-8674\(02\)01135-2](https://doi.org/10.1016/S0092-8674(02)01135-2)
- Wang J, Knaut H (2014) Chemokine signaling in development and disease. *Development* 141:4199-4205 . <https://doi.org/10.1242/dev.101071>
- Knorr M, Münzel T, Wenzel P (2014) Interplay of NK cells and monocytes in vascular inflammation and myocardial infarction. *Front Physiol* 5: . <https://doi.org/10.3389/fphys.2014.00295>
- De Filippo K, Dudeck A, Hasenberg M, Nye E, van Rooijen N, Hartmann K, Gunzer M, Roers A, Hogg N (2013) Mast cell and macrophage chemokines CXCL1/CXCL2 control the early stage of neutrophil recruitment during tissue inflammation. *Blood* 121:4930-4937 . <https://doi.org/10.1182/blood-2013-02-486217>
- Tamagawa-Mineoka R, Katoh N, Kishimoto S (2009) Platelets play important roles in the late phase of the immediate hypersensitivity reaction. *J Allergy Clin Immunol* 123:581-587.e9 . <https://doi.org/10.1016/j.jaci.2008.12.1114>
- Tamagawa-Mineoka R (2015) Important roles of platelets as immune cells in the skin. *J Dermatol Sci* 77:93-101 . <https://doi.org/10.1016/j.jdermsci.2014.10.003>
- Kunkel EJ, Kim CH, Lazarus NH, Vierra MA, Soler D, Bowman EP, Butcher EC (2003) CCR10 expression is a common feature of circulating and mucosal epithelial tissue IgA Ab-secreting cells. *J Clin Invest* 111:1001-1010 . <https://doi.org/10.1172/JCI17244>
- Wilson E, Butcher EC (2004) CCL28 Controls Immunoglobulin (Ig)A Plasma Cell Accumulation in the Lactating Mammary Gland and IgA Antibody Transfer to the Neonate. *J Exp Med* 200:805-809 . <https://doi.org/10.1084/jem.20041069>
- Morteau O, Gerard C, Lu B, Ghiran S, Rits M, Fujiwara Y, Law Y, Distelhorst K, Nielsen EM, Hill ED, Kwan R, Lazarus NH, Butcher EC, Wilson E (2008) An indispensable role for the chemokine receptor CCR10 in IgA antibody-secreting cell accumulation. *J Immunol Baltim Md* 181:6309-6315 . <https://doi.org/10.4049/jimmunol.181.9.6309>
- LIU G-X, LAN J, SUN Y, HU Y-J, JIANG G-S (2012) Expression of the chemokine CCL28 in pleomorphic adenoma and adenolymphoma of the human salivary glands. *Exp Ther Med* 4:65-69 . <https://doi.org/10.3892/etm.2012.544>
- Vazquez MI, Catalan-Dibene J, Zlotnik A (2015) B cells responses and cytokine production are regulated by their immune microenvironment. *Cytokine* 74:318-326 . <https://doi.org/10.1016/j.cyto.2015.02.007>
- Simon CMFrcp (2013) Underwood's Pathology: a Clinical Approach, 6 edition. Churchill Livingstone, Edinburgh
- Vinay KMMFrcp, KAM Abul, CAMP Jon (2017) Robbins Basic Pathology, 10 edition. Elsevier, Philadelphia, Pennsylvania
- Gillitzer R, Goebeler M (2001) Chemokines in cutaneous wound healing. *J Leukoc Biol* 69:513-521 . <https://doi.org/10.1189/jlb.69.4.513>
- Ishida Y, Gao J-L, Murphy PM (2008) Chemokine Receptor CXCR1 Mediates Skin Wound Healing by Promoting Macrophage and Fibroblast Accumulation and Function. *J Immunol* 180:569-579 . <https://doi.org/10.4049/jimmunol.180.1.569>
- Devalaraja RM, Nanney LB, Qian Q, Du J, Yu Y, Devalaraja MN, Richmond A (2000) Delayed Wound Healing in CXCR2 Knockout Mice. *J Invest Dermatol* 115:234-244 . <https://doi.org/10.1046/j.1523-1747.2000.00034.x>
- Michel G, Kemény L, Peter RU, Beetz A, Ried C, Arenberger P, Ruzicka T (1992) Interleukin-8 receptor-mediated chemotaxis of normal human epidermal cells. *FEBS Lett* 305:241-243 . [https://doi.org/10.1016/0014-5793\(92\)80677-9](https://doi.org/10.1016/0014-5793(92)80677-9)
- Tuschil A, Lam C, Haslberger A, Lindley I (1992) Interleukin-8 stimulates calcium transients and promotes epidermal cell proliferation. *J Invest Dermatol* 99:294-298 . <https://doi.org/10.1111/1523-1747.ep12616634>
- Gyulai R, Hunyadi J, Kenderessy-Szabó A, Kemény L, Dobozy A (1994) Chemotaxis of freshly separated and cultured human keratinocytes. *Clin Exp Dermatol* 19:309-311 . <https://doi.org/10.1111/j.1365-2230.1994.tb01201.x>
- Rennekampff HO, Hansbrough JF, Kiessig V, Doré C, Sticherling M, Schröder JM (2000) Bioactive interleukin-8 is expressed in wounds and enhances wound healing. *J Surg Res* 93:41-54 . <https://doi.org/10.1006/jsre.2000.5892>
- Niu G, Wright KL, Huang M, Song L, Haura E, Turkson J, Zhang S, Wang T, Sinibaldi D, Coppola D, Heller R, Ellis LM, Karras J, Bromberg J, Pardoll D, Jove R, Yu H (2002) Constitutive Stat3 activity up-regulates VEGF

- expression and tumor angiogenesis. *Oncogene* 21:2000-8 <https://doi.org/10.1038/sj.onc.1205260>
30. Ogata H, Sekikawa A, Yamagishi H, Ichikawa K, Tomita S, Imura J, Ito Y, Fujita M, Tsubaki M, Kato H, Fujimori T, Fukui H (2010) GRO α promotes invasion of colorectal cancer cells. *Oncol Rep* 24:1479-86. https://doi.org/10.3892/or_00001008
 31. Miyake M, Lawton A, Goodison S, Urquidí V, Gomes-Giacoaia E, Zhang G, Ross S, Kim J, Rosser CJ (2013) Chemokine (C-X-C) ligand 1 (CXCL1) protein expression is increased in aggressive bladder cancers. *BMC Cancer* 13:322. <https://doi.org/10.1186/1471-2407-13-322>
 32. Quintás-Cardama A, Verstovsek S (2013) Molecular pathways: Jak/STAT pathway: mutations, inhibitors, and resistance. *Clin Cancer Res* 19:1933-40. <https://doi.org/10.1158/1078-0432.CCR-12-0284>
 33. Katoh H, Wang D, Daikoku T, Sun H, Dey SK, Dubois RN (2013) CXCR2-expressing myeloid-derived suppressor cells are essential to promote colitis-associated tumorigenesis. *Cancer Cell* 24:631-44. <https://doi.org/10.1016/j.ccr.2013.10.009>
 34. Wei ZW, Xia GK, Wu Y, Chen W, Xiang Z, Schwarz RE, Brekken RA, Awasthi N, He YL, Zhang CH (2015) CXCL1 promotes tumor growth through VEGF pathway activation and is associated with inferior survival in gastric cancer. *Cancer Lett* 359:335-43. <https://doi.org/10.1016/j.canlet.2015.01.033>
 35. Xu J, Zhang C, He Y, Wu H, Wang Z, Song W, Li W, He W, Cai S, Zhan W (2012) Lymphatic endothelial cell-secreted CXCL1 stimulates lymphangiogenesis and metastasis of gastric cancer. *Int J Cancer* 130:787-97. <https://doi.org/10.1002/ijc.26035>
 36. Wang D, Yang W, Du J, Devalaraja MN, Liang P, Matsumoto K, Tsubakimoto K, Endo T, Richmond A (2000) MGSA/GRO-mediated melanocyte transformation involves induction of Ras expression. *Oncogene* 19:4647-59. <https://doi.org/10.1038/sj.onc.1203820>
 37. Yang G, Rosen DG, Zhang Z, Bast RC, Mills GB, Colacino JA, Mercado-Uribe I, Liu J (2006) The chemokine growth-regulated oncogene 1 (Gro-1) links RAS signaling to the senescence of stromal fibroblasts and ovarian tumorigenesis. *Proc Natl Acad Sci U S A* 103:16472-7. <https://doi.org/10.1073/pnas.0605752103>
 38. Dong YL, Kabir SM, Lee ES, Son DS (2013) CXCR2-driven ovarian cancer progression involves upregulation of proinflammatory chemokines by potentiating NF- κ B activation via EGFR-transactivated Akt signaling. *PLoS One* 8:e83789. <https://doi.org/10.1371/journal.pone.0083789>
 39. Ijichi H, Chytil A, Gorska AE, Aakre ME, Brier B, Tada M, Mohri D, Miyabayashi K, Asaoka Y, Maeda S, Ikenoue T, Tateishi K, Wright CV, Koike K, Omata M, Moses HL (2011) Inhibiting Cxcr2 disrupts tumor-stromal interactions and improves survival in a mouse model of pancreatic ductal adenocarcinoma. *J Clin Invest* 121:4106-17. <https://doi.org/10.1172/JCI42754>
 40. Oue E, Lee JW, Sakamoto K, Iimura T, Aoki K, Kayamori K, Michi Y, Yamashiro M, Harada K, Amagasa T, Yamaguchi A (2012) CXCL2 synthesized by oral squamous cell carcinoma is involved in cancer-associated bone destruction. *Biochem Biophys Res Commun* 424:456-61. <https://doi.org/10.1016/j.bbrc.2012.06.132>
 41. Hardaway AL, Herroon MK, Rajagurubandara E, Podgorski I (2015) Marrow adipocyte-derived CXCL1 and CXCL2 contribute to osteolysis in metastatic prostate cancer. *Clin Exp Metastasis*. <https://doi.org/10.1007/s10585-015-9714-5>
 42. Bachmeier B, Nerlich AG, Iancu CM, Cilli M, Schleicher E, Vené R, Dell'Eva R, Jochum M, Albini A, Pfeffer U (2007) The chemopreventive polyphenol Curcumin prevents hematogenous breast cancer metastases in immunodeficient mice. *Cell Physiol Biochem* 19:137-52. <https://doi.org/10.1159/000099202>
 43. Killian PH, Kronski E, Michalik KM, Barbieri O, Astigiano S, Sommerhoff CP, Pfeffer U, Nerlich AG, Bachmeier BE (2012) Curcumin inhibits prostate cancer metastasis in vivo by targeting the inflammatory cytokines CXCL1 and -2. *Carcinogenesis* 33:2507-19. <https://doi.org/10.1093/carcin/bgs312>
 44. Kronski E, Fiori ME, Barbieri O, Astigiano S, Mirisola V, Killian PH, Bruno A, Pagani A, Rovera F, Pfeffer U, Sommerhoff CP, Noonan DM, Nerlich AG, Fontana L, Bachmeier BE (2014) miR181b is induced by the chemopreventive polyphenol curcumin and inhibits breast cancer metastasis via down-regulation of the inflammatory cytokines CXCL1 and -2. *Mol Oncol* 8:581-95. <https://doi.org/10.1016/j.molonc.2014.01.005>
 45. Owen JL, Criscitiello MF, Libreros S, Garcia-Areas R, Guthrie K, Torroella-Kouri M, Iragavarapu-Charyulu V (2011) Expression of the inflammatory chemokines CCL2, CCL5 and CXCL2 and the receptors CCR1-3 and CXCR2 in T lymphocytes from mammary tumor-bearing mice. *Cell Immunol* 270:172-82. <https://doi.org/10.1016/j.cellimm.2011.05.004>
 46. Gardai SJ, McPhillips KA, Frasch SC, Janssen WJ, Starefeldt A, Murphy-Ullrich JE, Bratton DL, Oldenborg PA, Michalak M, Henson PM (2005) Cell-surface calreticulin initiates clearance of viable or apoptotic cells through trans-activation of LRP on the phagocyte. *Cell* 123:321-34. <https://doi.org/10.1016/j.cell.2005.08.032>
 47. Obeid M, Tesniere A, Ghiringhelli F, Fimia GM, Apetoh L, Perfettini JL, Castedo M, Mignot G, Panaretakis T, Casares N, Métivier D, Larochette N, van Endert P, Ciccosanti F, Piacentini M, Zitvogel L, Kroemer G (2007) Calreticulin exposure dictates the immunogenicity of cancer cell death. *Nat Med* 13:54-61. <https://doi.org/10.1038/nm1523>
 48. Sukkurwala AQ, Martins I, Wang Y, Schlemmer F, Ruckenstein C, Durchschlag M, Michaud M, Senovilla L, Sistigu A, Ma Y, Vacchelli E, Sulpice E, Gidrol X, Zitvogel L, Madeo F, Galluzzi L, Kepp O, Kroemer G (2014) Immunogenic calreticulin exposure occurs through a phylogenetically conserved stress pathway involving the chemokine CXCL8. *Cell Death Differ* 21:59-68. <https://doi.org/10.1038/cdd.2013.73>
 49. Gao JL, Murphy PM (1994) Human cytomegalovirus open reading frame US28 encodes a functional beta chemokine receptor. *J Biol Chem* 269:28539-42. [https://doi.org/10.1016/S0021-9258\(19\)61936-8](https://doi.org/10.1016/S0021-9258(19)61936-8)
 50. Damon I, Murphy PM, Moss B (1998) Broad spectrum chemokine antagonistic activity of a human poxvirus chemokine homolog. *Proc Natl Acad Sci U S A* 95:6403-7. <https://doi.org/10.1073/pnas.95.11.6403>
 51. Chen S, Bacon KB, Li L, Garcia GE, Xia Y, Lo D, Thompson DA, Siani MA, Yamamoto T, Harrison JK, Feng L (1998) In vivo inhibition of CC and CX3C chemokine-induced leukocyte infiltration and attenuation of glomerulonephritis in Wistar-Kyoto (WKY) rats by vMIP-II. *J Exp Med* 188:193-8. <https://doi.org/10.1084/jem.188.1.193>
 52. Vink C, Beisser PS, Bruggeman CA (1999) Molecular mimicry by cytomegaloviruses. Function of cytomegalovirus-encoded homologues of G protein-coupled receptors, MHC class I heavy chains and chemokines. *Intervirology* 42:342-9. <https://doi.org/10.1159/000053970>
 53. Milne RS, Mattick C, Nicholson L, Devaraj P, Alami A, Gompels UA (2000) RANTES binding and down-regulation by a novel human herpesvirus-6 beta chemokine receptor. *J Immunol* 164:2396-404. <https://doi.org/10.4049/jimmunol.164.5.2396>
 54. Kotwal GJ (2000) Poxviral mimicry of complement and chemokine system components: what's the end game? *Immunol Today* 21:242-8. [https://doi.org/10.1016/S0167-5699\(00\)01606-6](https://doi.org/10.1016/S0167-5699(00)01606-6)
 55. Murphy PM (2001) Viral exploitation and subversion of the immune system through chemokine mimicry. *Nat Immunol* 2:116-22. <https://doi.org/10.1038/84214>
 56. Tripp RA, Jones LP, Haynes LM, Zheng H, Murphy PM, Anderson LJ (2001) CX3C chemokine mimicry by respiratory syncytial virus G glycoprotein. *Nat Immunol* 2:732-8. <https://doi.org/10.1038/90675>
 57. Alami A (2003) Viral mimicry of cytokines, chemokines and their receptors. *Nat Rev Immunol* 3:36-50. <https://doi.org/10.1038/nri980>

58. Bryant NA, Davis-Poynter N, Vanderplassen A, Alcamí A (2003) Glycoprotein G isoforms from some alphaherpesviruses function as broad-spectrum chemokine binding proteins. *EMBO J* 22:833-46. <https://doi.org/10.1093/emboj/cdg092>
59. Lau EK, Allen S, Hsu AR, Handel TM (2004) Chemokine-receptor interactions: GPCRs, glycosaminoglycans and viral chemokine binding proteins. *Adv Protein Chem* 68:351-91. [https://doi.org/10.1016/S0065-3233\(04\)68010-7](https://doi.org/10.1016/S0065-3233(04)68010-7)
60. Rosenkilde MM (2005) Virus-encoded chemokine receptors--putative novel antiviral drug targets. *Neuropharmacology* 48:1-13. <https://doi.org/10.1016/j.neuropharm.2004.09.017>
61. Epperson ML, Lee CA, Fremont DH (2012) Subversion of cytokine networks by virally encoded decoy receptors. *Immunol Rev* 250:199-215. <https://doi.org/10.1111/imr.12009>
62. McSharry BP, Avdic S, Slobedman B (2012) Human cytomegalovirus encoded homologs of cytokines, chemokines and their receptors: roles in immunomodulation. *Viruses* 4:2448-70. <https://doi.org/10.3390/v4112448>
63. Wu L, Gerard NP, Wyatt R, Choe H, Parolin C, Ruffing N, Borsetti A, Cardoso AA, Desjardín E, Newman W, Gerard C, Sodroski J (1996) CD4-induced interaction of primary HIV-1 gp120 glycoproteins with the chemokine receptor CCR-5. *Nature* 384:179-183. <https://doi.org/10.1038/384179a0>
64. Liao F, Alkhatib G, Peden KWC, Sharma G, Berger EA, Farber JM (1997) STRL33, A Novel Chemokine Receptor-like Protein, Functions as a Fusion Cofactor for Both Macrophage-tropic and T Cell Line-tropic HIV-1. *J Exp Med* 185:2015-2023. <https://doi.org/10.1084/jem.185.11.2015>
65. Farber JM (1998) Chemokines, lymphocytes, and HIV. *Braz J Med Biol Res* 31:11-17. <https://doi.org/10.1590/S0100-879X1998000100002>
66. Rizzuto CD, Wyatt R, Hernández-Ramos N, Sun Y, Kwong PD, Hendrickson WA, Sodroski J (1998) A conserved HIV gp120 glycoprotein structure involved in chemokine receptor binding. *Science* 280:1949-1953. <https://doi.org/10.1126/science.280.5371.1949>
67. Wyatt R, Sodroski J (1998) The HIV-1 Envelope Glycoproteins: Fusogens, Antigens, and Immunogens. *Science* 280:1884-1888. <https://doi.org/10.1126/science.280.5371.1884>
68. Efremov R, Truong M-J, Darcissac ECA, Zeng J, Grau O, Vergoten G, Debarb C, Capron A, Bahr GM (1999) Human chemokine receptors CCR5, CCR3 and CCR2B share common polarity motif in the first extracellular loop with other human G-protein coupled receptors. *Eur J Biochem* 263:746-756. <https://doi.org/10.1046/j.1432-1327.1999.00553.x>
69. Rowland-Jones S (1999) The role of chemokine receptors in HIV infection. *Sex Transm Infect* 75:148-151. <https://doi.org/10.1136/sti.75.3.148>
70. Suresh P, Wanchu A (2006) Chemokines and chemokine receptors in HIV infection: role in pathogenesis and therapeutics. *J Postgrad Med* 52:210-217.
71. Nedellec R, Coetzer M, Shimizu N, Hoshino H, Polonis VR, Morris L, Mårtensson UEA, Binley J, Overbaugh J, Mosier DE (2009) Virus Entry via the Alternative Coreceptors CCR3 and FPRL1 Differs by Human Immunodeficiency Virus Type 1 Subtype. *J Virol* 83:8353-8363. <https://doi.org/10.1128/JVI.00780-09>
72. Wang Z, Shang H, Jiang Y (2017) Chemokines and Chemokine Receptors: Accomplices for Human Immunodeficiency Virus Infection and Latency. *Front Immunol* 8. <https://doi.org/10.3389/fimmu.2017.01274>
73. Choe H, Farzan M, Sun Y, Sullivan N, Rollins B, Ponath PD, Wu L, Mackay CR, LaRosa G, Newman W, Gerard N, Gerard C, Sodroski J (1996) The β -Chemokine Receptors CCR3 and CCR5 Facilitate Infection by Primary HIV-1 Isolates. *Cell* 85:1135-1148. [https://doi.org/10.1016/S0092-8674\(00\)81313-6](https://doi.org/10.1016/S0092-8674(00)81313-6)
74. Doranz BJ, Rucker J, Yi Y, Smyth RJ, Samson M, Peiper SC, Parmentier M, Collman RG, Doms RW (1996) A dual-tropic primary HIV-1 isolate that uses fusin and the beta-chemokine receptors CKR-5, CKR-3, and CKR-2b as fusion cofactors. *Cell* 85:1149-1158. [https://doi.org/10.1016/S0092-8674\(00\)81314-8](https://doi.org/10.1016/S0092-8674(00)81314-8)
75. Deng HK, Unutmaz D, KewalRamani VN, Littman DR (1997) Expression cloning of new receptors used by simian and human immunodeficiency viruses. *Nature* 388:296-300. <https://doi.org/10.1038/40894>
76. Louisiriratchanakul S, Liu H, Roongpisuthipong A, Nakayama E, Takebe Y, Shioda T (2002) Genetic Analysis of HIV-1 Discordant Couples in Thailand: Association of CCR2 64I Homozygosity With HIV-1-Negative Status. *J Acquir Immune Defic Syndr* 1999 29:314-5. <https://doi.org/10.1097/00042560-200203010-00014>
77. Furci L, Scarlatti G, Burastero S, Tambussi G, Colognesi C, Quillent C, Longhi R, Loverro P, Borgonovo B, Gaffi D, Carrow E, Malnati M, Lusso P, Siccardi AG, Lazzarin A, Beretta A (1997) Antigen-driven C-C Chemokine-mediated HIV-1 Suppression by CD4+ T Cells from Exposed Uninfected Individuals Expressing the Wild-type CCR-5 Allele. *J Exp Med* 186:455-460. <https://doi.org/10.1084/jem.186.3.455>
78. Nedoszytko B, sokolowska-wojdylo M, Ruckemann-Dziurdzińska K, Roszkiewicz J, Nowicki R (2014) Chemokines and cytokines network in the pathogenesis of the inflammatory skin diseases: Atopic dermatitis, psoriasis and skin mastocytosis. *Postępy Dermatol Alergol* 31:84-91. <https://doi.org/10.5114/pdia.2014.40920>
79. Zhou X, Krueger JG, Kao M-CJ, Lee E, Du F, Menter A, Wong WH, Bowcock AM (2003) Novel mechanisms of T-cell and dendritic cell activation revealed by profiling of psoriasis on the 63,100-element oligonucleotide array. *Physiol Genomics* 13:69-78. <https://doi.org/10.1152/physiolgenomics.00157.2002>
80. Nomura I, Gao B, Boguniewicz M, Darst MA, Travers JB, Leung DYM (2003) Distinct patterns of gene expression in the skin lesions of atopic dermatitis and psoriasis: A gene microarray analysis. *J Allergy Clin Immunol* 112:1195-1202. <https://doi.org/10.1016/j.jaci.2003.08.049>
81. Martynowicz H, Janus A, Nowacki D, Mazur G (2014) The Role of Chemokines in Hypertension. *Adv Clin Exp Med Off Organ Wroclaw Med Univ* 23:319-325. <https://doi.org/10.17219/acem/37123>
82. Rudemiller NP, Crowley SD (2017) The role of chemokines in hypertension and consequent target organ damage. *Pharmacol Res* 119:404-411. <https://doi.org/10.1016/j.phrs.2017.02.026>
83. Tanase DM, Gosav EM, Radu S, Ouatu A, Rezus C, Ciocoiu M, Costea CF, Floria M (2019) Arterial Hypertension and Interleukins: Potential Therapeutic Target or Future Diagnostic Marker? In: *Int. J. Hypertens.* <https://www.hindawi.com/journals/ijhy/2019/3159283/>. Accessed 25 Sep 2020. <https://doi.org/10.1155/2019/3159283>
84. Chistiakov DA, Orekhov AN, Sobenin IA, Bobryshev YV (2014) Plasmacytoid dendritic cells: development, functions, and role in atherosclerotic inflammation. *Front Physiol* 5. <https://doi.org/10.3389/fphys.2014.00279>
85. Marx N, Mach F, Sauty A, Leung JH, Sarafi MN, Ransohoff RM, Libby P, Plutzky J, Luster AD (2000) Peroxisome Proliferator-Activated Receptor- γ Activators Inhibit IFN- γ -Induced Expression of the T Cell-Active CXC Chemokines IP-10, Mig, and I-TAC in Human Endothelial Cells. *J Immunol Baltim Md* 1950 164:6503-6508. <https://doi.org/10.4049/jimmunol.164.12.6503>
86. Rotondi M, Chiovato L, Romagnani S, Serio M, Romagnani P (2007) Role of Chemokines in Endocrine Autoimmune Diseases. *Endocr Rev* 28:492-520. <https://doi.org/10.1210/er.2006-0044>
87. Kimura H, Caturegli P (2007) Chemokine orchestration of autoimmune thyroiditis. *Thyroid Off J Am Thyroid Assoc* 17:1005-1011. <https://doi.org/10.1089/thy.2007.0267>
88. Ferrari SM, Fallahi P, Vita R, Antonelli A, Benvenega S (2015) Peroxisome Proliferator-Activated Receptor- γ in Thyroid Autoimmunity. In: *PPAR*

- Res. <https://www.hindawi.com/journals/ppar/2015/232818/>. Accessed 25 Sep 2020. <https://doi.org/10.1155/2015/232818>
89. Ferrari SM, Ruffilli I, Elia G, Ragusa F, Paparo SR, Patrizio A, Mazzi V, Antonelli A, Fallahi P (2019) Chemokines in hyperthyroidism. *J Clin Transl Endocrinol* 16: . <https://doi.org/10.1016/j.jcte.2019.100196>
 90. Domberg J, Chao L, Papewalis C, Pfleger C, Xu K, Willenberg HS, Hermesen D, Scherbaum WA, Schlöot NC, Schott M (2008) Circulating Chemokines in Patients with Autoimmune Thyroid Diseases. *Horm Metab Res* 40:416-421. <https://doi.org/10.1055/s-2008-1073151>
 91. Liu C, Cui G, Zhu M, Kang X, Guo H (2014) Neuroinflammation in Alzheimer's disease: chemokines produced by astrocytes and chemokine receptors. *Int J Clin Exp Pathol* 7:8342-8355
 92. Alsadany MA, Shehata HH, Mohamad MI, Mahfouz RG (2013) Histone Deacetylases Enzyme, Copper, and IL-8 Levels in Patients With Alzheimer's Disease. *Am J Alzheimers Dis Dementiasr* 28:54-61 . <https://doi.org/10.1177/1533317512467680>
 93. Galimberti D, Schoonenboom N, Scheltens P, Fenoglio C, Venturelli E, Pijnenburg Y a. L, Bresolin N, Scarpini E (2006) Intrathecal chemokine levels in Alzheimer disease and frontotemporal lobar degeneration. *Neurology* 66:146-147 . <https://doi.org/10.1212/01.wnl.0000191324.08289.9d>
 94. Venturelli E, Galimberti D, Fenoglio C, Lovati C, Finazzi D, Guidi I, Corrà B, Scalabrini D, Clerici F, Mariani C, Forloni G, Bresolin N, Scarpini E (2006) Candidate gene analysis of IP-10 gene in patients with Alzheimer's disease. *Neurosci Lett* 404:217-221 . <https://doi.org/10.1016/j.neulet.2006.05.054>
 95. Tripathy D, Thirumangalakudi L, Grammas P (2010) RANTES upregulation in the Alzheimer's disease brain: a possible neuroprotective role. *Neurobiol Aging* 31:8-16 . <https://doi.org/10.1016/j.neurobiolaging.2008.03.009>
 96. Kester MI, van der Flier WM, Visser A, Blankenstein MA, Scheltens P, Oudejans CB (2011) Decreased mRNA expression of CCL5 [RANTES] in Alzheimer's disease blood samples. *Clin Chem Lab Med* 50:61-65. <https://doi.org/10.1515/CCLM.2011.731>
 97. Laske C, Stellos K, Stransky E, Seizer P, Akcay O, Eschweiler GW, Leyhe T, Gawaz M (2008) Decreased plasma and cerebrospinal fluid levels of stem cell factor in patients with early Alzheimer's disease. *J Alzheimers Dis JAD* 15:451-460 . <https://doi.org/10.3233/jad-2008-15311>
 98. van der Meer P, Ulrich AM, González-Scarano F, Lavi E (2000) Immunohistochemical analysis of CCR2, CCR3, CCR5, and CXCR4 in the human brain: potential mechanisms for HIV dementia. *Exp Mol Pathol* 69:192-201 . <https://doi.org/10.1006/exmp.2000.2336>
 99. Williams JL, Holman DW, Klein RS (2014) Chemokines in the balance: maintenance of homeostasis and protection at CNS barriers. *Front Cell Neurosci* 8: . <https://doi.org/10.3389/fncel.2014.00154>
 100. Deverman BE, Patterson PH (2009) Cytokines and CNS Development. *Neuron* 64:61-78. <https://doi.org/10.1016/j.neuron.2009.09.002>
 101. Harrison JK, Jiang Y, Chen S, Xia Y, Maciejewski D, McNamara RK, Streit WJ, Salafranca MN, Adhikari S, Thompson DA, Botti P, Bacon KB, Feng L (1998) Role for neuronally derived fractalkine in mediating interactions between neurons and CX3CR1-expressing microglia. *Proc Natl Acad Sci* 95:10896-10901. <https://doi.org/10.1073/pnas.95.18.10896>
 102. Huising MO, Meulen T van der, Flik G, Kemenade BMLV (2004) Three novel carp CXC chemokines are expressed early in ontogeny and at nonimmune sites. *Eur J Biochem* 271:4094-4106 . <https://doi.org/10.1111/j.1432-1033.2004.04347.x>
 103. Banisadr G, Bhattacharyya B, Belmadani A, Izen SC, Ren D, Tran PB, Miller RJ (2011) The chemokine BRAK/CXCL14 regulates synaptic transmission in the adult mouse dentate gyrus stem cell niche. *J Neurochem* 119:1173-1182 . <https://doi.org/10.1111/j.1471-4159.2011.07509.x>
 104. Kokovay E, Goderie S, Wang Y, Lotz S, Lin G, Sun Y, Roysam B, Shen Q, Temple S (2010) Adult SVZ lineage cells home to and leave the vascular niche via differential responses to SDF1/CXCR4 signaling. *Cell Stem Cell* 7:163-173 . <https://doi.org/10.1016/j.stem.2010.05.019>
 105. Fan Q, Gayen M, Singh N, Gao F, He W, Hu X, Tsai L-H, Yan R (2019) The intracellular domain of CX3CL1 regulates adult neurogenesis and Alzheimer's amyloid pathology. *J Exp Med* 216:1891-1903 . <https://doi.org/10.1084/jem.20182238>
 106. Sahin H, Trautwein C, Wasmuth HE (2010) Functional role of chemokines in liver disease models. *Nat Rev Gastroenterol Hepatol* 7:682-690 . <https://doi.org/10.1038/nrgastro.2010.168>
 107. Saiman Y, Friedman S (2012) The Role of Chemokines in Acute Liver Injury. *Front Physiol* 3:213 . <https://doi.org/10.3389/fphys.2012.00213>
 108. Marra F, Tacke F (2014) Roles for Chemokines in Liver Disease. *Gastroenterology* 147:577-594.e1. <https://doi.org/10.1053/j.gastro.2014.06.043>
 109. Roehlen N, Crouchet E, Baumert TF (2020) Liver Fibrosis: Mechanistic Concepts and Therapeutic Perspectives. *Cells* 9: . <https://doi.org/10.3390/cells9040875>
 110. Oo YH, Shetty S, Adams DH (2010) The Role of Chemokines in the Recruitment of Lymphocytes to the Liver. *Dig Dis* 28:31-44 . <https://doi.org/10.1159/000282062>
 111. Bhatia M, Zemans RL, Jeyaseelan S (2012) Role of Chemokines in the Pathogenesis of Acute Lung Injury. *Am J Respir Cell Mol Biol* 46:566-572 . <https://doi.org/10.1165/rcmb.2011-0392TR>
 112. Chen H, Wang D, Wang X (2010) Role of Airway Epithelium-Origin Chemokines and their Receptors in COPD. *J Epithel Biol Pharmacol* 3:
 113. Barnes PJ (2016) Inflammatory mechanisms in patients with chronic obstructive pulmonary disease. *J Allergy Clin Immunol* 138:16-27 . <https://doi.org/10.1016/j.jaci.2016.05.011>
 114. Henrot P, Prevel R, Berger P, Dupin I (2019) Chemokines in COPD: From Implication to Therapeutic Use. *Int J Mol Sci* 20: . <https://doi.org/10.3390/ijms20112785>
 115. Keane MP (2008) The role of chemokines and cytokines in lung fibrosis. *Eur Respir Rev* 17:151-156 . <https://doi.org/10.1183/09059180.00010908>
 116. Blease K, Lukacs NW, Hogaboam CM, Kunkel SL (2000) Chemokines and their role in airway hyper-reactivity. *Respir Res* 1:54-61 . <https://doi.org/10.1186/rr13>
 117. El Mesallamy HO, Ahmed HH, Bassyouni AA, Ahmed AS (2012) Clinical significance of inflammatory and fibrogenic cytokines in diabetic nephropathy. *Clin Biochem* 45:646-650 . <https://doi.org/10.1016/j.clinbiochem.2012.02.021>
 118. Gaballa M, Farag Y (2012) Predictors of diabetic nephropathy. *Cent Eur J Med* 8: . <https://doi.org/10.2478/s11536-012-0055-3>
 119. Lim AKH, Tesch GH (2012) Inflammation in Diabetic Nephropathy. In: *Mediators Inflamm.* <https://www.hindawi.com/journals/mi/2012/146154/>. Accessed 25 Sep 2020. <https://doi.org/10.1155/2012/146154>
 120. Sanchez AP, Sharma K (2009) Transcription factors in the pathogenesis of diabetic nephropathy. *Expert Rev Mol Med* 11:e13 . <https://doi.org/10.1017/S1462399409001057>
 121. Takebayashi K, Matsumoto S, Aso Y, Inukai T (2006) Association between circulating monocyte chemoattractant protein-1 and urinary albumin excretion in nonobese Type 2 diabetic patients. *J Diabetes Complications* 20:98-104 . <https://doi.org/10.1016/j.jdiacomp.2005.05.008>

122. Belkaid Y, Hand TW (2014) Role of the Microbiota in Immunity and Inflammation. *Cell* 157:121-141 .
<https://doi.org/10.1016/j.cell.2014.03.011>
123. Garlet GP, Sfeir CS, Little SR (2014) Restoring Host-Microbe Homeostasis via Selective Chemoattraction of Tregs. *J Dent Res* 93:834-839 . <https://doi.org/10.1177/0022034514544300>
124. Zheng D, Liwinski T, Elinav E (2020) Interaction between microbiota and immunity in health and disease. *Cell Res* 30:492-506 .
<https://doi.org/10.1038/s41422-020-0332-7>
125. Park UC, Kim TW, Yu HG (2014) Immunopathogenesis of Ocular Behçet's Disease. In: *J. Immunol. Res.* <https://www.hindawi.com/journals/jir/2014/653539/>. Accessed 13 Sep 2020. <https://doi.org/10.1155/2014/653539>
126. Frassanito MA, Dammacco R, Cafforio P, Dammacco F (1999) Th1 polarization of the immune response in Behçet's disease: a putative pathogenetic role of interleukin-12. *Arthritis Rheum* 42:1967-1974 .
[https://doi.org/10.1002/1529-0131\(199909\)42:9<1967::AID-ANR24>3.0.CO;2-Z](https://doi.org/10.1002/1529-0131(199909)42:9<1967::AID-ANR24>3.0.CO;2-Z).
127. Houman H, Hamzaoui A, Ben Ghorbal I, Khanfir M, Feki M, Hamzaoui K (2004) Abnormal expression of chemokine receptors in Behçet's disease: relationship to intracellular Th1/Th2 cytokines and to clinical manifestations. *J Autoimmun* 23:267-273 .
<https://doi.org/10.1016/j.jaut.2004.07.005>
128. Ahmed MB, Houman H, Miled M, Dellagi K, Louzir H (2004) Involvement of chemokines and Th1 cytokines in the pathogenesis of mucocutaneous lesions of Behçet's disease. *Arthritis Rheum* 50:2291-2295 . <https://doi.org/10.1002/art.20334>
129. Imamura Y, Kurokawa MS, Yoshikawa H, Nara K, Takada E, Masuda C, Tsukikawa S, Ozaki S, Matsuda T, Suzuki N (2005) Involvement of Th1 cells and heat shock protein 60 in the pathogenesis of intestinal Behçet's disease. *Clin Exp Immunol* 139:371-378 . <https://doi.org/10.1111/j.1365-2249.2005.02695.x>
130. Dalghous AM, Freysdottir J, Fortune F (2006) Expression of cytokines, chemokines, and chemokine receptors in oral ulcers of patients with Behçet's disease (BD) and recurrent aphthous stomatitis is Th1-associated, although Th2-association is also observed in patients with BD. *Scand J Rheumatol* 35:472-475 .
<https://doi.org/10.1080/03009740600905380>