

12-12-2017

## ANTIOXIDANT TRACE ELEMENTS AND OXIDATIVE STRESS STATUS IN DAIRY CATTLE SUFFERING from MASTITIS

Medhat Nassif

*Department of Animal Medicine, Faculty of Veterinary Medicine, Kafr El-sheikh University, Kafr El-sheikh, Egypt*

Hussam Ibrahim

*Department of Internal Medicine, Infectious Diseases and Fish Diseases, Faculty of Veterinary Medicine, Mansoura University, Mansoura, Egypt*

Heba Moharam

*Department of Internal Medicine, Infectious Diseases and Fish Diseases, Faculty of Veterinary Medicine, Mansoura University, Mansoura, Egypt*

Follow this and additional works at: <https://mvmj.researchcommons.org/home>

---

### How to Cite This Article

Nassif, Medhat; Ibrahim, Hussam; and Moharam, Heba (2017) "ANTIOXIDANT TRACE ELEMENTS AND OXIDATIVE STRESS STATUS IN DAIRY CATTLE SUFFERING from MASTITIS," *Mansoura Veterinary Medical Journal*: Vol. 18: Iss. 1, Article 7.

DOI: <https://doi.org/10.21608/mvmj.2017.126425>

This Original Article is brought to you for free and open access by Mansoura Veterinary Medical Journal. It has been accepted for inclusion in Mansoura Veterinary Medical Journal by an authorized editor of Mansoura Veterinary Medical Journal.

---

## Mansoura Veterinary Medical Journal

---

# ANTIOXIDANT TRACE ELEMENTS AND OXIDATIVE STRESS STATUS IN DAIRY CATTLE SUFFERING FROM MASTITIS

Medhat Nassif Nassif Mohamed\*, Hussam Mohamed Mohamed Ibrahim\*,  
Heba Adel Moharam Mahmoud\*\*

*\*Department of Animal Medicine, Faculty of Veterinary Medicine, Kafr El-sheikh*

*University, Kafr El-sheikh, Egypt*

*\*\*Department of Internal Medicine, Infectious Diseases and Fish Diseases, Faculty of Veterinary Medicine,  
Mansoura University, Mansoura, Egypt*

---

### ABSTRACT

*Mastitis is an inflammation of the mammary glands caused by several pathogens. Reactive oxygen species are natural end products of the intensive cellular metabolism. Oxidative stress occurs when there is a disturbance of homeostasis causing inflammation of the mammary gland. This study was performed to evaluate blood antioxidant profile and trace elements in dairy cows with clinical mastitis. For this aim, venous blood samples were collected from 116 cows (100 mastitic, 16 control). In cows with clinical mastitis, there was a significant ( $p < 0.05$ ) increase in the clinical index score compared with control group. Biochemically, there was a significant ( $p < 0.05$ ) decrease in super oxide dismutase, catalase and total antioxidant capacity as well as in the level of zinc. However, there was a significant ( $p < 0.05$ ) increase in the levels of malondialdehyde and glutathione reductase as well as in that of copper. The results of the current study indicate that the body antioxidant defense system was compromised in dairy cows with clinical mastitis creating a state of oxidative stress.*

---

### INTRODUCTION

Mastitis may be defined as an inflammation of all structures forming the mammary tissue and the surrounding connective tissue. The disease is the reaction of the mammary gland to irritants and significantly influences the quality and quantity of mammary tissue and milk (Yüksel *et al.*, 2009). It is mainly resulting in pathological, physical and chemical changes in milk and glandular tissues, resulting from an invasion of mammary tissues by pathogenes through the

teat canal as mentioned by Quinn *et al.*, (2002) and Radostits *et al.*, (2007). The disease results in some changes in the composition of milk and shortening of the productive lives of cows (Gürbulak *et al.*, 2009). It is important to improve productivity and quality of milk by reducing mastitis incidence (Halasa *et al.*, 2007; Cha *et al.*, 2011 and Hogeveen *et al.*, 2011).

Mastitis was found to induce economic losses including value of discarded milk as milk from affected cows may become unsuitable for human consumption due to

---

bacterial contamination resulting in food poisoning, interference with manufacturing process and in rare cases providing mechanism of spread of zoonotic diseases among population. Zoonotic diseases transmitted via raw cow milk included Staphylococcal food poisoning, brucellosis, tuberculosis, toxoplasmosis and leptospirosis as previously stated by **Mungube *et al.*, (2005)** and **Radostits *et al.*, (2007)**.

Oxidative stress is usually described as an imbalance between antioxidant and oxidant levels as mentioned by **Lykkesfeldt and Svendsen, (2007)**. A condition of oxidative stress was produced when the production of oxidants exceeded the capacity of antioxidant defense resulting in oxidative damage to macromolecules such as lipid, proteins and DNA (**Sordillo and Aitken, 2009**).

In this regard, **Jin *et al.*, (2014)** mentioned that Mammary epithelial cells exhibit a high metabolic rate during lactation and therefore large amounts of lipid peroxides and reactive oxygen species (ROS) are produced in vivo.

A few hours after the udder infection with pathogenic microorganisms, the number of somatic cells in the milk, was found to be increased in response to activation of the inflammatory processes. When the mammary gland was invaded and was colonized by bacteria, the macrophages responded by initiating the inflammatory response, one that attracted polymorphonuclear cells in milk to kill the bacteria. More than 90% of SCC in infected glands was composed of neutrophils (**Groza, 2006; Andrieu, 2008**). More over **Rinaldi *et al.*, (2007)** stated that Reactive oxygen species (ROS) mediated the antibacterial activity of neutrophils.

An excess of oxidative reactions of bacterial processes might cause damage to the tissues. The absence of optimal amounts of antioxidants and an excess of ROS resulted in oxidative stress development (**Andrei, 2010b**).

Free radicals are natural end products of the intensive metabolism in cells of the living organism, including high-yielding dairy cows. When the homeostasis was disturbed mainly by generation and accumulation of these free radicals, oxidative processes resulted in oxidative stress causing mastitis in dairy cows (**Polawska *et al.*, 2012**).

The objective of the present study was to evaluate the role of oxidative stress biomarkers, antioxidant enzymes and metabolites as possible biomarkers for clinical mastitis in dairy cows.

## MATERIALS AND METHODS

A total number of 116 dairy cows aged between three to ten years were included in this study. Of all, 100 cows exhibited the clinical signs of mastitis. Besides, 16 apparently healthy cows were selected as a control group. All samples were collected in the period between October, 2014 and January, 2015, where 96 samples were collected from Dakahlia while 20 samples were collected from Damietta Governorate, Egypt.

According to **Radostits *et al.*, (2007)**, detailed clinical examination of the animals was carried out. Physical examination of the cows was concerned with evidence of systemic illness (on which fever, ruminal stasis, inappetence to anorexia, tachycardia, dullness and recumbency was observed and recorded). Also, local physical examination of the udder via palpation and inspection of the quarters as well as the supramammary lymph nodes was conducted. It was directed toward detection of the inflammatory cardinal signs or evidence of fibrosis or atrophy. For observation of any abnormalities in the examined milk, visual examination of the milk was done including (milk flakes, clots, pus, watery and bloody secretion).

## Sampling:

### 1. Milk sample

Under complete aseptic precautions, sampling of the examined milk was taken using standard methods described by **Hogan et al. (1999)**. About (20 ml) of milk was obtained from each affected quarter in separate sterile and well corked falkon tubes. The obtained milk samples were incubated at 37 °c for 12-24 hours, then centrifugate the tubes at 3000 rpm for 15 minutes for concentration of the bacterial cells at the sediment and discard the supernatant. From the sediment, a loopful of about (0.01ml) was mixed with 5 ml of TryptoneSoya Broth (**Oxoid, LTD, Batingstock, Hampshire, England**) in sterile corked test tubes. Then, the tubes were incubated for 18 hours. Each sample was streaked on three different selective media. For isolation of *Staphylococcus aureus*, Baird parker (Bp.) medium (**Oxoid**) supplemented with 5% egg yolk- telluriteemulsion was used. Edward's medium (EDW.) (**Oxoid**) supplemented with 6% defibrinated sheep blood was used for the selective isolation of *Streptococcus spp.* For *Escherichiacoli* selective isolation, Eosin methelene blue medium (EMB.) was used (**Quinn et al., 2002**).

### 2. Blood Samples Collection

Two types of venous blood samples (ten ml for each) were collected via puncture of jugular vein from each cow. **Blood plasma samples**, were collected into Eppendorf tube which was mixed with Ethylenediaminetetraacetic acid (EDTA) as anticoagulant for biochemical estimation of level and activity of SOD, CAT and GR. hematological studies. **Blood serum sample**,

were collected in clean dry tube without anti-coagulant. The collected blood samples were left to coagulate, then centrifuged at 3000 rpm for 10 minutes to obtain blood serum for biochemical analysis of TAC, MDA, Zn and Cu levels according to the method described by manufacture (**Biogiagnostic, Cairo, Egypt**).

### 3. Statistical analysis

Data were subjected to statistical analysis using statistical software program (SPSS for Windows, version 15, USA). Means and standard deviation for each variable were estimated. Differences between means of different groups were carried out using one way ANOVA with Duncan multiple comparison tests. Correlations between different parameters were carried out using Pearson correlation coefficient. Differences between means at  $p < 0.05$  were considered significant.

## RESULTS

Concerning the bacteriological isolation and identification, *Staphylococcus aureus* appeared as black shiny colonies surround by hallow zone on Baird parker. *Escherichia coli* (*E. coli*) appeared as green metallic sheen colonies on the Eosin Methelene Blue media. While *Streptococcus agalactiae* (*S. agalactiae*) appeared on Edward's media (B hemolysis) as colorless colonies with bluish hue and surrounded with complete zone of hemolysis. In the current study, the bacteriological isolation profile showed that about (159) bacterial isolates were obtained from the (100) cultured milk samples. The mixed infection 118/159 (74.21%) between *Staphylococcus spp.*, *Streptococcus spp.* and *E.coli* was the

largest percentage of isolation, then isolation percentage, came to be for *Staphylococcus spp.* 21/159 (13.22%) then for *E. coli* 15/159 (9.43%) and *Streptococcus spp.* 5/159 (3.14%).

Regarding the level of different oxidative stress markers and enzymatic antioxidants activities in clinical mastitis, there was a significant ( $p < 0.05$ ) decrease in Super oxide dismutase (SOD), Catalase (CAT) and Total antioxidant capacity (TAC) compared to those

of the control group. Whereas there was a significant ( $p < 0.05$ ) increase in Malondialdehyde (MDA) and Glutathione reductase (GR) compared to control group.

Regarding different antioxidant trace elements levels in clinical mastitis, there was a significant ( $p < 0.05$ ) decrease in Zn level compared to control group. Whereas there was a significant ( $p < 0.05$ ) increase in Cu level compared to those of the control group.

**Table 1:** Clinical signs of cows affected with mastitis caused by *Staphylococcal spp.*, *E. coli*, *Streptococcal spp.* and mixed infection.

Clinical variables	Staphylococcus spp.	Streptococcus spp.	E. coli	Mixed infection
	Number of animal affected(n=21)	Number of animal affected(n=5)	Number of animal affected(n=15)	Number of animal affected(n=59)
<b>Evidences of fever with systemic illness</b>				
Mild(39.5°C- 39.9°C)	16	4	2	38
Sever(40°C- 41.4°C)	5	1	13	21
No systemic illness	0	0	0	0
<b>Udder abnormalities</b>				
Cardinal signs of inflammation	21	5	15	59
Evidence of fibrosis or atrophy	0	0	0	0
<b>physical characters of milk</b>				
Pus and greenish discoloration	2	2	1	6
Yellowish discoloration	5	0	10	16
Milk clots	7	3	4	30
Milk tinged with blood	5	0	0	6
Whole bloody milk	2	0	0	1

**Table 2:** Different Oxidative Stress Markers, Enzymatic Antioxidants Activity and Trace elements level (mean values  $\pm$  SD) in Clinically Healthy cows and in those with clinical mastitis disease.

Groups	SOD	CAT	MDA	TAC	GR	Zn	Cu
	(U/mL)	(U/L)	(nmol/mL)	(mM/L)	(U/L)	( $\mu$ g/dl)	( $\mu$ g/dl)
<b>Control</b> (n = 16)	219.8 $\pm$ 50.8 <sup>a</sup>	494.6 $\pm$ 147.0 <sup>a</sup>	15.1 $\pm$ 2.1 <sup>a</sup>	1.1 $\pm$ 0.2 <sup>a</sup>	934.5 $\pm$ 306.6 <sup>a</sup>	267.4 $\pm$ 64.4 <sup>a</sup>	70.2 $\pm$ 7.2 <sup>a</sup>
<b>E-coli</b> (n=15)	151.7 $\pm$ 36.0 <sup>b</sup>	406.8 $\pm$ 122.8 <sup>a</sup>	17.7 $\pm$ 2.2 <sup>b</sup>	0.9 $\pm$ 0.1 <sup>b</sup>	1214.9 $\pm$ 448.0 <sup>b</sup>	190.5 $\pm$ 9.0 <sup>b</sup>	77.2 $\pm$ 2.3 <sup>b</sup>
<b>Staphylococcus</b> (n=21)	159.5 $\pm$ 37.5 <sup>b</sup>	349.8 $\pm$ 126.8 <sup>b</sup>	16.4 $\pm$ 1.4 <sup>a</sup>	0.9 $\pm$ 0.1 <sup>b</sup>	1298.4 $\pm$ 335.2 <sup>b</sup>	193.0 $\pm$ 13.3 <sup>b</sup>	76.8 $\pm$ 16.8 <sup>b</sup>
<b>Streptococcus</b> (n = 5)	164.5 $\pm$ 39.0 <sup>b</sup>	242.3 $\pm$ 207.4 <sup>b</sup>	17 $\pm$ 0.8 <sup>b</sup>	0.8 $\pm$ 0.1 <sup>b</sup>	1171.6 $\pm$ 256.3 <sup>ab</sup>	184.9 $\pm$ 3.6 <sup>b</sup>	77.0 $\pm$ 1.8 <sup>b</sup>
<b>Mixed</b> (n = 59)	157.6 $\pm$ 35.6 <sup>b</sup>	294.8 $\pm$ 147.9 <sup>c</sup>	16.9 $\pm$ 1.9 <sup>b</sup>	0.9 $\pm$ 0.1 <sup>b</sup>	1153.8 $\pm$ 274 <sup>ab</sup>	190.6 $\pm$ 11.4 <sup>b</sup>	76.5 $\pm$ 2.7 <sup>b</sup>

## DISCUSSION

All clinically examined dairy cows, in the current study, displayed a significant increase in the clinical index score. These findings were in agreement with those reported by **Biffa et al. (2005)** and **ENREF\_38Bedane et al. (2012)**. Such findings might be attributed to the inflammation of the mammary glands following their infection which represents the stage at which clinical mastitis occurs with appearance of gross abnormalities of the milk, variable degrees of clinical abnormalities of the udder and variable systemic effects. Moreover, the clinical signs of the disease was found to be varied in severity depending on the

causative microorganisms and their virulence (**Radostits et al., 2007**).

In the present study SOD was significantly decreased compared to those of the control group. These results were similar to those reported by **Machado et al., (2014)** who attributed such decrease to the low level of zinc in the blood of mastitic cows.

In the current study, there was a significant decrease in catalase activity which was similar to other finding observed by **Jhambh et al. (2013)**. The decreased enzymatic antioxidant activities might be related to the increased consumption to counteract the effect of reactive oxygen species as catalase is an enzyme that catalyzes the decomposition of hydrogen peroxide into hydroxyl radicals and water.

Lipid peroxidation was a non-enzymatic chain reaction led to creation of lipid peroxides and other intermediates, based on oxidation of mainly unsaturated fatty acids. These intermediates including malondialdehyde (MDA) might influence the properties of cell membranes (Catalá, 2010). In the present study, malondialdehyde was significantly increased comparatively to the control group. These results were similar to those reported by Jhambh *et al.*, (2013). During clinical mastitis, the enhanced concentrations of plasma MDA, could be due to excessive ROS production from the clinically inflamed mammary gland suggesting a compromise in antioxidant defense of the body (Boulanger *et al.*, 2002).

In the current study, the antioxidant defense system was compromised in dairy cows with clinical mastitis, which was evidenced by decreased plasma TAC and increased MDA level, which might indirectly indicate increased whole activity of free radical reflecting the worse state of oxidative stress in such cases (Celi, 2010).

Glutathione reductase was significantly increased comparatively to the control group. The involvement of glutathione dependent enzymes (glutathione peroxidases and glutathione reductase) would be important for lipid peroxides neutralization during clinical mastitis. These results were similar to those previously reported by (Halliwell and Chirico, 1993). Also our findings may be owing to the enhancement of the coupled activities of the glutathione dependent enzymes, leading to intense regeneration of reduced glutathione (GSH) from oxidized glutathione (GSSH) obtained after reduction of peroxides into alcohols as previously stated by Kizil *et al.* (2007).

Trace elements as copper and zinc played an important role in several biochemical processes as they are essential component in the antioxidant enzymes as SOD and CAT (Yang and Li, 2014). In the current study zinc was significantly decreased comparatively to the control group. Such results were similar to the result previously reported by Ranjan *et al.*, (2005). Its deficiency was associated with decreased leukocyte function and increased susceptibility to bacterial infection (Hayajneh, 2014) with a resultant of a state of oxidative stress (Zago and Oteiza, 2001). Moreover, the decreased serum zinc level was regarded as non-specific host defense mechanism against bacterial infection, which would reduce the availability of that divalent cation needed for bacterial growth as previously stated by (Failla, 2003).

Copper is involved in the antioxidant system via its involvement in ceruloplasmin and the enzymes Cu-Zn SOD. Ceruloplasmin is a Cu transport protein that also exhibits oxidase activity. In addition it is an acute phase protein that increases during infection and may be important in scavenging superoxide radicals (Broadley and Hoover, 1989). In the present study copper was significantly increased compared to control group, these results were similar to those recorded by (Ranjan *et al.*, 2005). Ceruloplasmin contains more than 95% of the circulating copper in animals. Its level in blood has been reported to increase many fold during inflammation. Therefore, increase in blood copper level in the clinical cases of mastitis might be due to elevated level of ceruloplasmin in inflammatory udder conditions (Conner *et al.*, 1986).

## REFERENCES

- Andrei, S. (2010b).** Correlations Between Antioxidant Enzymes Activity and Lipids Peroxidation Level in Blood and Milk from Cows with Subclinical Mastitis. *Bulletin of University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca. Veterinary Medicine*, 67, 6-11.
- Andrieu, S. (2008).** Is there a role for organic trace element supplements in transition cow health? *The Veterinary Journal*, 176, 77-83.
- Bedane, A., Kasim, G., Yohannis, T., et al. (2012).** Study on prevalence and risk factors of bovine mastitis in Borana Pastoral and agro-pastoral settings of Yabello District, Borana Zone, Southern Ethiopia. *American-Eurasian Journal of Agriculture and Environmental Science*, 12, 1274-1281.
- Biffa, D., Debela, E., Beyene, F., et al. (2005).** Factors associated with bovine udder infections in smallholder dairy farms in Ethiopia  
Facteurs liés aux infections de la mamelle bovine dans les petites exploitations laitières en Ethiopie. *Bulletin of Animal Health and Production in Africa*, 53, 258-265.
- Boulanger, V., Zhao, X. & Lacasse, P. (2002).** Protective Effect of Melatonin and Catalase in Bovine Neutrophil-induced Model of Mammary Cell Damage. *Journal of Dairy Science*, 85, 562-569.
- Broadley, C. & Hoover, R. L. (1989).** Ceruloplasmin reduces the adhesion and scavenges superoxide during the interaction of activated polymorphonuclear leukocytes with endothelial cells. *The American journal of pathology*, 135, 647.
- Catalá, A. (2010).** A synopsis of the process of lipid peroxidation since the discovery of the essential fatty acids. *Biochemical and Biophysical Research Communications*, 399, 318-323.
- Celi, P. (2010).** Biomarkers of oxidative stress in ruminant medicine. *Immunopharmacology and Immunotoxicology*, 33, 233-240.
- Cha, E., Bar, D., Hertl, J. A., et al. (2011).** The cost and management of different types of clinical mastitis in dairy cows estimated by dynamic programming. *Journal of Dairy Science*, 94, 4476-4487.
- Conner, J., Eckersall, P., Doherty, M., et al. (1986).** Acute phase response and mastitis in the cow. *Research in veterinary science*, 41, 126.
- Failla, M. L. (2003).** Trace elements and host defense: recent advances and continuing challenges. *The Journal of nutrition*, 133, 1443S-1447S.
- Groza, I. (2006).** Ginecologie, andrologie și obstetrică veterinară—Compendiu. Editura Academiei Române, București.
- Gürbulak, K., Canoğlu, E., Abay, M., et al. (2009).** Determination of Subclinical Mastitis in Dairy Cows by Different Methods. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi*, 15, 765-770.
- Halasa, T., Huijps, K., Østerås, O., et al. (2007).** Economic effects of bovine mastitis and mastitis management: A review. *Veterinary Quarterly*, 29, 18-31.



- Halliwell, B. & Chirico, S. (1993).** Lipid peroxidation: its mechanism, measurement, and significance. *The American journal of clinical nutrition*, 57, 715S-724S.
- Hayajneh, F. M. F. (2014).** Plasma ascorbic acid levels in sheep infected with hydated cyst.
- Hogan, J., Gonzalez, R. & Harmon, R. (1999).** Laboratory Handbook on Bovine Mastitis. Revised ed. Madison, Wisconsin: Nat Mastitis Counc. Inc.
- Hogeveen, H., Huijps, K. & Lam, T. (2011).** Economic aspects of mastitis: New developments. *New Zealand Veterinary Journal*, 59, 16-23.
- Jhambh, R., Dimri, U., Gupta, V., et al. (2013).** Blood antioxidant profile and lipid peroxides in dairy cows with clinical mastitis. *Vet World*, 6, 271.
- Jin, L., Yan, S., Shi, B., et al. (2014).** Effects of vitamin A on the milk performance, antioxidant functions and immune functions of dairy cows. *Animal Feed Science and Technology*, 192, 15-23.
- Kizil, O., Akar, Y., Saat, N., et al. (2007).** The plasma lipid peroxidation intensity (MDA) and chain-breaking antioxidant concentrations in the cows with clinic or subclinic mastitis. *Revue Med Vet*, 158, 529-533.
- Lykkesfeldt, J. & Svendsen, O. (2007).** Oxidants and antioxidants in disease: Oxidative stress in farm animals. *The Veterinary Journal*, 173, 502-511.
- Machado, V. S., Oikonomou, G., Lima, S. F., et al. (2014).** The effect of injectable trace minerals (selenium, copper, zinc, and manganese) on peripheral blood leukocyte activity and serum superoxide dismutase activity of lactating Holstein cows. *The Veterinary Journal*, 200, 299-304.
- Mungube, E., Tenhagen, B.-A., Regassa, F., et al. (2005).** Reduced milk production in udder quarters with subclinical mastitis and associated economic losses in crossbred dairy cows in Ethiopia. *Tropical animal health and production*, 37, 503-512.
- Polawska, E., Bagnicka, A. W., Niemczuk, K., et al. (2012).** Relations between the oxidative status, mastitis, milk quality and disorders of reproductive functions in dairy cows—a review. *Animal Science Papers and Reports*, 30, 297-307.
- Quinn, P. J., Carter, M. E., Markey, B., et al. (2002).** Clinical Veterinary Microbiology Microbial diseases, bacterial causes of bovine mastitis, 8.
- Radostits, O., Gay, C. C., Hinchcliff, K., et al. (2007).** Veterinary medicine: a textbook of the diseases of cattle, sheep, pigs, goats and horses. 10, 1045-1046.
- Ranjan, R., Swarup, D., Naresh, R., et al. (2005).** Enhanced Erythrocytic Lipid Peroxides and Reduced Plasma Ascorbic Acid, and Alteration in Blood Trace Elements Level in Dairy Cows with Mastitis. *Veterinary Research Communications*, 29, 27-34.
- Rinaldi, M., Moroni, P., Paape, M. J., et al. (2007).** Evaluation of assays for the measurement of bovine neutrophil reactive oxygen species. *Veterinary Immunology and Immunopathology*, 115, 107-125.

- Sordillo, L. M. & Aitken, S. L. (2009).** Impact of oxidative stress on the health and immune function of dairy cattle. *Veterinary Immunology and Immunopathology*, 128, 104-109.
- Yang, F. L. & Li, X. S. (2014).** Role of antioxidant vitamins and trace elements in mastitis in dairy cows. *Journal of Advanced Veterinary and Animal Research*, 2, 1-9.
- Yüksel, M., Kandemir, F., Deveci, H., et al. (2009).** The serum ALP, ALT and glucose levels of healthy and subclinically mastitic cows. *Atatürk Üniversitesi Veteriner Bilimleri Dergisi*, 4, 163-168.
- Zago, M. P. & Oteiza, P. I. (2001).** The antioxidant properties of zinc: interactions with iron and antioxidants. *Free Radical Biology and Medicine*, 31, 266-274.

## الملخص العربي

### العناصر النزرة المضادة للأكسدة وحالة الأكسدة في الأبقار الحلوب المصابة بالتهاب الضرع

مدحت ناصف ناصف محمد<sup>\*</sup>، حسام محمد محمد إبراهيم<sup>\*\*</sup>، هبة عادل محرم محمود<sup>\*\*</sup>

<sup>\*</sup> قسم طب الحيوان- كلية الطب البيطري- جامعة كفر الشيخ- كفر الشيخ- مصر

<sup>\*\*</sup> قسم أمراض الباطنة والمعدية والأسماك- كلية الطب البيطري- جامعة المنصورة- المنصورة- مصر

مرض التهاب الضرع هو التهاب في الغدد الثدييه الذي يسببه العديد من المسببات المرضية. تعتبر أنواع الأكسجين النشطة هي الناتج الطبيعي النهائي للأيض الخلوي. يحدث الإجهاد التأكسدي عندما يكون هناك اضطراب في التوازن ويؤدي ذلك إلي التهاب الضرع في الأبقار عالية الإنتاج. وقد أجريت هذه الدراسة لتقييم دور مضادات الأكسدة في دم الأبقار التي تعاني من التهاب الضرع.

لقد تم تجميع عينات الدم الوريدي عشوائيا من ١٦ بقرة لاتعاني من أمراض كمجموعة ضابطة و ١٠٠ بقرة حلوب مصابة بالتهاب الضرع بناءا علي الفحص الطبي الشامل. وجد في الأبقار المصابة بالتهاب الضرع أن هناك زيادة ملحوظة في مؤشر الفحص الإكلينيكي مقارنة بالمجموعة الضابطة.

لوحظ أن هناك نقص نوعي في قيمة السوبر أكسيد ديسموتيز و الكاتاليز و القدرة المضادة للأكسدة الكلية والزنك بينما كانت هناك زيادة ملحوظة في المألون داي الديهيد و الجلوتاثيون ريداكناز و النحاس.

نستخلص من نتائج الدراسة الحالية أن هناك تأثير سلبي علي نظام الدفاع المضاد للأكسدة بالجسم في الأبقار المصابة بالتهاب الضرع الإكلينيكي مؤدياً الي حالة من الإجهاد التأكسدي.